Chronic Pain Scale for Nonverbal Adults With Intellectual Disabilities (CPS-NAID)

Please indicate how often this person has shown the signs referred to in *items 1-24* in the <u>last 5 minutes</u>.

Please circle a number for each item. If an item does not apply to this person (for example, this person cannot reach with his/her hands), then indicate "not applicable" for that item.

0 =	Not present at all during the observation period. (Note if the item is not present because the person is not capable of performing that act, it should be scored as "NA").
1 =	Seen or heard rarely (hardly at all), but is present.
2 =	Seen or heard a number of times, but not continuous (not all the time).
3 =	Seen or heard often, almost continuous (almost all the time); anyone would easily notice this if they saw the person for a few moments during the observation time.
NA =	Not applicable. This person is not capable of performing this action.

0 = Not at all 1 = Just a little 2 = Fairly Often 3 = Ver		y Often	NA = Not Applicable					
Moaning, whining, whimpering (fairly soft)					1	2	3	NA
2. Crying (moderately loud)					1	2	3	NA
3. A specific sound or word for pain (e.g. A word, cry or type of laugh)					1	2	3	NA
4. Not cooperating, irritable, unhappy					1	2	3	NA
5. Less interaction with others, withdrawn					1	2	3	NA
6. Seeking comfort of physical closeness					1	2	3	NA
7. Being difficult to distract, not able to satisfy or pacify					1	2	3	NA
8. A furrowed brow					1	2	3	NA
9. A change in eyes, including: squinching of eyes opened wide, eyes					1	2	3	NA
frowning								
10. Turning down	0	1	2	3	NA			
	11. Lips puckering up, tight, pouting or quivering					2	3	NA
		g or thrusting tongue	out	0	1	2	3	NA
13. Not moving, less active, quiet					1	2	3	NA
14. Stiff, spastic, te	0	1	2	3	NA			
	r touching part of the			0	1 1	2	3	NA
	16. Protecting, favouring or guarding part of body that hurts					2	3	NA
17. Flinching or moving the body part away, being sensitive to touch					1	2	3	NA
18. Moving the body in a specific way to show pain (e.g. Head back, arms down, curls up, etc.)					1	2	3	NA
19. Shivering	0	1	2	3	NA			
20. Change in colour, pallor					1	2	3	NA
21. Sweating, perspiring					1	2	3	NA
22. Tears					1	2	3	NA
23. Sharp intake of breath, gasping					1	2	3	NA
24. Breath holding					1	2	3	NA
Subtotals:								
1. For each subtotal write the number of times each value was chosen					1x	2 x	3x	NA
2. Multiply the value of each selection by how many times that value was chosen								Total:
3. Add each subtotal to find the total score					=	=	=	

SCORING:

- 1. Add up the scores for each item to compute the Total Score. Items marked "NA" are scored as "0" (zero).
- 2. Check whether the score is greater than the cut-off score.

A score of <u>10 or greater</u> means that there is a 94% chance that the person <u>has pain</u>.

A score of 9 or lower means that there is an 87% chance that the person does not have pain.